

Possible probe of the QCD odderon singularity through the quasidiffractive η_c production in $\gamma\gamma$ collisions .

L. MOTYKA^a, J. KWIECIŃSKI^b,

^a*Institute of Physics, Jagellonian University, Cracow, Poland*

^b*Department of Theoretical Physics,*

H. Niewodniczański Institute of Nuclear Physics, Cracow, Poland

Abstract

The reactions $\gamma\gamma \rightarrow \eta_c\eta_c$ and $\gamma\gamma \rightarrow \eta_c + X$ are discussed within the three gluon exchange model. We give predictions for the differential cross-sections and discuss feasibility of measuring these processes at LEP2 and TESLA. The total cross-sections were estimated to be approximately equal to 40 fb and 120 fb for $\gamma\gamma \rightarrow \eta_c\eta_c$ and $\gamma\gamma \rightarrow \eta_c + X$ respectively assuming exchange of elementary gluons that corresponds to the odderon with intercept equal to unity. These values can be enhanced by a factor equal to 1.9 and 2.1 for LEP2 and TESLA energies if the odderon intercept is equal to 1.07. The estimate of cross-sections $\sigma(e^+e^- \rightarrow e^+e^-\eta_c\eta_c)$ and $\sigma(e^+e^- \rightarrow e^+e^-\eta_cX)$ for untagged e^+ and e^- is also given.

The dominant contribution in the high-energy limit of perturbative QCD is given by the exchange of interacting gluons [1, 2]. The fact that the gluons have spin equal to unity automatically implies that the cross-sections corresponding to the exchange of elementary gluons are independent of the incident energies while the interaction between gluons leads to increasing cross-section. Besides the pomeron [1, 2, 3, 4, 5] one also expects presence of the so called “odderon” singularity [1, 2, 3, 6, 7]. In the leading logarithmic approximation the pomeron is described by the BFKL equation which corresponds to the sum of ladder diagrams with the reggeized gluon exchange along the ladder. The odderon is described by the three-gluon exchange. Unlike pomeron which corresponds to the vacuum quantum numbers and so to the positive charge conjugation the odderon is characterised by $C = -1$ (and $I=0$) i.e. it carries the same quantum numbers as the ω Regge pole. The (phenomenologically) determined intercept λ_ω of the ω Regge pole is approximately equal to $1/2$ [8]. The novel feature of the odderon singularity corresponding to the gluonic degrees of freedom is the potentially very high value of its intercept $\lambda_{odd} \gg \lambda_\omega$. The exchange of the three (noninteracting) gluons alone generates singularity with intercept equal to unity while interaction between gluons described in the leading logarithmic approximation by the BKP equation [6, 7, 9] is even capable to boost the odderon intercept above unity [10]¹. The energy dependence of the amplitudes corresponding to $C = -1$ exchange becomes similar to the diffractive ones which are controlled by the pomeron exchange.

Possible tests of both QCD perturbative pomeron as well as of odderon have to rely on (semi) hard processes where presence of hard scale can justify the use of perturbative QCD. Very useful measurement in this respect is the very high energy exclusive photo (or electro) production of heavy charmonia (i.e. J/Ψ [12, 13] or η_c [14, 15, 16] etc. for probing the QCD pomeron or odderon respectively). The estimate of the odderon contribution to the photo- (or electro-) production of even charge conjugation mesons does however require model assumptions about the coupling of the three gluon system to a proton. It would therefore be useful to consider the process which could in principle be calculated entirely within perturbative QCD. The relevant measurement which fulfills those criteria is the exclusive quasidiffractive production of even C charmonia in $\gamma\gamma$ collisions or, to be precise, the processes: $\gamma\gamma \rightarrow \eta_c\eta_c$ or $\gamma\gamma \rightarrow \eta_c + X$ [17, 18]. The main purpose of our paper is to present the theoretical and phenomenological description of the double η_c production in high energy $\gamma\gamma$ collisions and of the process $\gamma\gamma \rightarrow \eta_c + X$ with the large rapidity gap between η_c and the hadronic system X assuming the three gluon exchange mechanism. In our paper we shall follow the formalism developed in ref [18] where the production of pseudoscalar mesons in $\gamma\gamma$ collisions within the three

¹The results of ref. [10] are however in conflict with recent analysis [11]




Figure 1: Kinematics of the three gluon exchange mechanism of the process $\gamma\gamma \rightarrow \eta_c\eta_c$ (1a) and $\gamma\gamma \rightarrow \eta_c + X$ (1b).

gluon exchange mechanism is discussed with great detail.

The kinematics of the three gluon exchange diagram to the processes: $\gamma^*(q_1) + \gamma^*(q_2) \rightarrow \eta_c\eta_c$ and $\gamma^*(q_1) + \gamma^*(q_2) \rightarrow \eta_c + X$ is illustrated in Fig. 1a and Fig. 1b.

The amplitude M^{ij} for the process $\gamma^*(q_1) + \gamma^*(q_2) \rightarrow \eta_c\eta_c$ which corresponds to the transverse polarisation of both photons can be written in the following way:

$$M^{ij} = W^2 C_{qq} \int \frac{d^2\delta_1 d^2\delta_2}{\delta_1^2 \delta_2^2 \delta_3^2} \Phi_{\gamma\eta_c}^i(Q_1^2, \delta_1, \delta_2, \Delta) \Phi_{\gamma\eta_c}^j(Q_2^2, -\delta_1, -\delta_2, -\Delta) \quad (1)$$

where i and j are the polarisation indices and C_{qq} is given by:

$$C_{qq} = \frac{10}{9\pi N_c^2} \alpha_s^3(m_c^2) \quad (2)$$

and Δ and δ_i denote the transverse components of Δ and δ_i while $Q_i^2 = -q_i^2$.

The relevant diagrams describing the impact factor $\Phi_{\gamma\eta_c}^{i,j}$ are given in Fig. 2. Besides the diagrams presented in Fig. 2 one has also to include those with the reversed direction of the quark lines. In the nonrelativistic approximation one gets the following expression for $\Phi_{\gamma\eta_c}^i$:

$$\Phi_{\gamma\eta_c}^i = F_{\eta_c} \sum_{k=1}^2 \epsilon_{ik} \left[\frac{\Delta^k}{\bar{M}_1^2 + \Delta^2} + \sum_{s=1}^3 \frac{2\delta_s^k - \Delta^k}{\bar{M}_1^2 + (2\delta_s - \Delta)^2} \right] \quad (3)$$




Figure 2: The diagrams defining the impact factor $\Phi_{\gamma\eta_c}$.

where

$$F_{\eta_c} = \sqrt{\frac{4m_{\eta_c}\Gamma_{\eta_c\rightarrow\gamma\gamma}}{\alpha q_c^2}} \quad (4)$$

and

$$\bar{M}_1^2 = 4m_c^2 + Q_1^2. \quad (5)$$

In equations (3,4,5) $m_c, m_{\eta_c}, \Gamma_{\eta_c\rightarrow\gamma\gamma}$ and q_c denote the charmed quark mass, the mass of the η_c , the η_c radiative width and the charge of the charm quark respectively. The formula for the impact factor $\Phi_{\gamma\eta_c}^j$ corresponding to the lower vertex is given by equation (3) after changing the polarisation index i into j and after reversing the sign of δ_s and of Δ in this equation and after changing $\bar{M}_1^2$ into $\bar{M}_2^2$ given by equation (5) with Q_2^2 instead of Q_1^2 . The approximations leading to the formula (3) are discussed in [15].

For $Q_1^2 = Q_2^2 \simeq 0$ it is convenient to represent the amplitude M^{ij} in the following way:

$$M^{ij} = W^2 \left(M_1 g^{ij} + M_2 \frac{\Delta^i \Delta^j}{\Delta^2} \right) \quad (6)$$

The corresponding formula for the differential cross-section averaged over the transverse




Figure 3: The diagrams defining the impact factor $\Phi_{\gamma X}$.

photons polarizations for the process $\gamma\gamma \rightarrow \eta_c \eta_c$ reads:

$$\frac{d\sigma}{dt} = \frac{1}{64\pi} [(M_1 + M_2)^2 + M_1^2] \quad (7)$$

For real $\gamma - s$ we have, of course, set $Q_{1,2}^2 = 0$.

The process $\gamma\gamma \rightarrow \eta_c + X$ is given by the diagram of Fig. 1b and the amplitude which corresponds to this diagram can be written as below:

$$M_{\gamma\gamma \rightarrow \eta_c + X} = W^2 \tilde{C}_{qq} \int \frac{d^2\delta_1 d^2\delta_2}{\delta_1^2 \delta_2^2 \delta_3^2} \Phi_{\gamma\eta_c}(Q_2^2, -\delta_1, -\delta_2, -\Delta) \Phi_{\gamma X}(Q_1^2, \delta_1, \delta_2, \Delta, p_1, p_2) \quad (8)$$

where

$$\Phi_{\gamma\eta_c} = \sum_{i,j=1}^2 \epsilon_2^i \epsilon_{ij} \Phi_{\gamma\eta_c}^j \quad (9)$$

and

$$\tilde{C}_{qq} = \frac{10}{9\pi N_c^2} [\alpha_s(m_c^2) \tilde{\alpha}_s]^{3/2}. \quad (10)$$

The coupling constant $\tilde{\alpha}_s$ is defined by the hard scale characteristic for the upper vertex in Fig. 1b. The impact factor $\Phi_{\gamma X}$ is given by the following equation:

$$\Phi_{\gamma X}(Q_1^2, \delta_1, \delta_2, \Delta, p_1, p_2) = -ieq_f \bar{u}(p_1) [m_f R \not{\epsilon}_1 + 2z \bar{Q} \epsilon_1 + \bar{Q} \not{\epsilon}_1] \not{q}_2 v(p_2) \quad (11)$$

where

$$\bar{\mathbf{Q}} = \left[\frac{\mathbf{p}_1}{m_f^2 + \mathbf{p}_1^2} + \sum_{s=1}^3 \frac{\boldsymbol{\delta}_s - \mathbf{p}_1}{m_f^2 + (\boldsymbol{\delta}_s - \mathbf{p}_1)^2} \right] + (\mathbf{p}_1 \rightarrow \mathbf{p}_2) \quad (12)$$

and

$$R = \left[\frac{-1}{m_f^2 + \mathbf{p}_1^2} + \sum_{s=1}^3 \frac{1}{m_f^2 + (\boldsymbol{\delta}_s - \mathbf{p}_1)^2} \right] + (\mathbf{p}_1 \rightarrow \mathbf{p}_2). \quad (13)$$

In equations (11, 12,13) $\bar{\mathbf{Q}}$ is a four-vector with transverse components $\bar{\mathbf{Q}}$ and vanishing longitudinal componenets, m_f denotes the mass of the (light) quark produced as the pair of $q\bar{q}$ jets, q_f its charge while z is the component of the photon four momentum q_1 carried by a quark jet. The four momenta p_1 and p_2 denote the four momenta of the quark (antiquark) in the final state and $\mathbf{p}_1, \mathbf{p}_2$ are their transverse parts respectively. When calculating the cross-sections we shall make the "equivalent quark approximation" [18, 19] which corresponds to the approximation of setting m_f equal to zero and to retaining the dominant term in $\bar{\mathbf{Q}}$ i.e.:

$$\bar{\mathbf{Q}} = \frac{\mathbf{p}_1}{\mathbf{p}_1^2} + \frac{\mathbf{p}_2}{\mathbf{p}_2^2}. \quad (14)$$

The remaining terms in (12) can also be large for $\boldsymbol{\delta}_s \simeq \mathbf{p}_i$ but their dependence on $\boldsymbol{\delta}_s$ is such that after the integration over $\boldsymbol{\delta}_s$ performed in (8) they are suppressed in comparison to the leading terms.

The differential cross-section for the process $\gamma\gamma \rightarrow \eta_c + X(q\bar{q})$ averaged over photon polarisations is given by the following formula:

$$d\sigma = \frac{(2\pi)^4}{2W^2} \sum_{\{\dots\}} M_{\gamma\gamma \rightarrow \eta_c + X}^+ M_{\gamma\gamma \rightarrow \eta_c + X} dPS_3(\gamma\gamma \rightarrow \eta_c + X(q\bar{q})) \quad (15)$$

where $\{\dots\}$ stands for incident photons helicities, outgoing light quark colours, flavours and polarisations and $dPS_3(\gamma\gamma \rightarrow \eta_c + X(q\bar{q}))$ is the standard parametrisation of the Lorentz invariant tree-body phase space. The decomposition property

$$dPS_3(\gamma\gamma \rightarrow \eta_c + X(q\bar{q})) = (2\pi)^3 dM_X^2 dPS_2(\gamma\gamma \rightarrow \eta_c + X) dPS_2(X \rightarrow q\bar{q}) \quad (16)$$

is employed and integration over invariant mass squared M_X^2 of the $q\bar{q}$ system is performed. The remaining integration over the two body phase space $dPS_2(X \rightarrow q\bar{q})$ in the applied approximation can not be extended to the regions $\mathbf{p}_i^2 \simeq 0$ since the integral would then be divergent. We do therefore introduce, following ref. [18, 19] a physical cut-off μ which is defined by the light quarks constituent masses and we set for its magnitude $\mu = 0.3 \text{ GeV}$. The integral is bounded from above by the condition

$\mathbf{p}_i^2 < \Delta^2$ which assures the diffractive nature of the process. Thus we obtain a logarithmic expression $\ln |t|/\mu^2$ arising from the integral $\int_{\mu^2}^{|t|} d\mathbf{p}_1^2/\mathbf{p}_1^2$. Finally the differential cross-section reads

$$\frac{d\sigma}{dt} = \frac{1}{64\pi} \left| \tilde{C}_{qq} \int \frac{d^2\delta_1 d^2\delta_2}{\delta_1^2 \delta_2^2 \delta_3^2} \Phi_{\gamma\eta_c}(Q_2^2, -\delta_1, -\delta_2, -\Delta, p_1, p_2) \right|^2 \frac{2\alpha\bar{e}^2}{3\pi} \ln \frac{|t|}{\mu^2} \quad (17)$$

where $\bar{e}^2 = N_c(e_u^2 + e_d^2 + e_s^2) = 2$ and e_i are the charges of the light quarks u , d and s respectively.

In Fig. 4a we show the differential cross-section of the process $\gamma\gamma \rightarrow \eta_c\eta_c$. Unlike the photoproduction of η_c on a nucleon target this cross-section does not vanish at $t = 0$. The result of the theoretical calculations can be conveniently represented for $|t| < 15 \text{ GeV}^2$ by the exponential form:

$$\frac{d\sigma}{dt} = 6.6 \text{ fb/GeV}^2 \exp(0.167 \text{ GeV}^{-2} t). \quad (18)$$

For the integrated cross-section we get $\sigma_{\eta_c\eta_c}^{tot} = 43 \text{ fb}$. The differential cross section for the process $\gamma\gamma \rightarrow \eta_c + X$ is presented in Fig. 4b. For $|t| < 15 \text{ GeV}^2$ it can be fitted to the exponential form:

$$\frac{d\sigma}{dt} = 64 \text{ fb/GeV}^2 \exp(0.25 \text{ GeV}^{-2} t). \quad (19)$$

The integrated cross-section is now $\sigma_{\eta_c X}^{tot}(|t| > 3 \text{ GeV}^2) = 120 \text{ fb}$. In our calculation we set $\alpha_s(m_c^2) = 0.38$, $\tilde{\alpha}_s = 0.3$, $m_c = 1.4 \text{ GeV}$, $m_{\eta_c} = 2.98 \text{ GeV}$ and $\Gamma_{\eta_c \rightarrow \gamma\gamma} = 7 \text{ keV}$.

The calculated cross-sections are energy independent since they correspond to the exchange of three elementary and non-interacting gluons. In this approximation the odderon singularity has its intercept λ_{odd} equal to unity. Interaction between gluons can boost this intercept above unity and one can take approximately this effect into account by multiplying the cross-sections by the enhancement factor

$$A_{enh}(W^2) = \bar{x}^{2(1-\lambda_{odd})} \quad (20)$$

where

$$\bar{x} = \frac{M_{\eta_c}^2}{W^2}. \quad (21)$$

The $\gamma\gamma$ system in e^+e^- collisions has a continuous spectrum. The C.M. energy squared W^2 of the $\gamma\gamma$ system is:

$$W^2 = z_1 z_2 s \quad (22)$$




Figure 4: Differential cross sections of the processes $\gamma\gamma \rightarrow \eta_c \eta_c$ (4a) and $\gamma\gamma \rightarrow \eta_c + X$ (4b) for $Q_1^2 \simeq Q_2^2 \simeq 0 \text{ GeV}^2$.

where z_i are the energy fractions of electrons (positrons) carried by the exchanged photons and s denotes the C.M. energy squared of the e^+e^- system. The distribution of those energy fractions is given by the standard flux-factors $f_{\gamma/e}(z, Q_{min}, Q_{max})$ of the virtual photons. In the equivalent photon approximation it is given by the following formula [20]:

$$f_{\gamma/e}(z, Q_{min}, Q_{max}) = \frac{\alpha}{2\pi} \left[\frac{1 + (1-z)^2}{z} \ln \frac{Q_{max}^2}{Q_{min}^2} \right] \quad (23)$$

where we have neglected a small term proportional to the electron mass squared m_e^2 . Q_{min}^2 and Q_{max}^2 in eq. (23) denote the minimal and maximal values of the photon virtuality. For untagged experiments the former is given by the kinematical limit:

$$Q_{min}^2 = \frac{m_e^2 z}{1-z} \quad (24)$$

and the latter by the antitagging condition $\theta_{e^\pm} < \theta_{max}$ which gives

$$Q_{max}^2 = (1-z)E_{beam}^2 \theta_{max}^2 \quad (25)$$

where $\theta_{e^\pm}$ denotes the scattering angle of the scattered $e^\pm$. Following ref. [20] we set $\theta_{max} = 30 \text{ mrad}$. The cross-section for the process $e^+e^- \rightarrow e^+e^- + Y$ which for untagged $e^\pm$ corresponds to the production of the hadronic state Y in the collision of almost real (virtual) photons is given by the following convolution integral:

$$\sigma_{e^+e^- \rightarrow e^+e^- + Y} =$$

Table 1: The estimated cross-sections $\sigma(e^+e^- \rightarrow e^+e^-\eta_c\eta_c)$ and $\sigma(e^+e^- \rightarrow e^+e^-\eta_cX)$.

$\sqrt{s}$ [GeV]	$\lambda_{odd} - 1$	$\sigma(e^+e^- \rightarrow e^+e^-\eta_c\eta_c)$ [fb]	$\sigma(e^+e^- \rightarrow e^+e^-\eta_cX)$ [fb]
180	0	1.3	4
180	0.07	2.5	7
500	0	3.5	10
500	0.07	7.4	21

$$\int_0^1 dz_1 \int_0^1 dz_2 \Theta(W^2 - W_{Y0}^2) \sigma_{\gamma\gamma \rightarrow Y}(W^2) f_{\gamma/e}(z_1, Q_{min}, Q_{max}) f_{\gamma/e}(z_2, Q_{min}, Q_{max}). \quad (26)$$

In order to estimate the effective enhancement factor due to $\lambda_{odd} > 1$ we have compared the convolution integrals:

$$I(s, \lambda_{odd}) = \int_0^1 dz_1 \int_0^1 dz_2 \Theta(z_{max} - M_{\eta_c}^2/(z_1 z_2 s)) f_{\gamma/e}(z_1, Q_{min}, Q_{max}) f_{\gamma/e}(z_2, Q_{min}, Q_{max}) \bar{x}^{2(1-\lambda_{odd})} \quad (27)$$

for $\lambda_{odd} = 1$ with that calculated for $\lambda_{odd} = 1.07$. We set $z_{max} = 0.05$ so that the $\gamma\gamma$ system is in the high energy (i.e. Regge) region. The ratio $A = I(s, \lambda_{odd} = 1.07)/I(s, \lambda_{odd} = 1)$ should give the expected enhancement factor for the given value of s . We get $A = 1.9$ and $A = 2.1$ for the LEP2 and TESLA energies respectively. The relatively small change in A with increasing s is caused by the fact that the convolution integral is dominated by small values of z_i .

The magnitude of the estimated cross-sections $\sigma(e^+e^- \rightarrow e^+e^-\eta_c\eta_c)$ and $\sigma(e^+e^- \rightarrow e^+e^-\eta_cX)$ for untagged e^+ and e^- in the final state are summarized in Table 1. We give values of those cross sections for two different C.M. energies of incident leptons corresponding to LEP2 and TESLA energies and for two different values of the odderon intercept $\lambda_{odd} = 1$ and $\lambda_{odd} = 1.07$. It may be seen from this table that the cross-sections are very small and so it may in particular be difficult to measure them with presently available luminosity at LEP2 [20].

To sum up we have applied the formalism developed in ref.[18] to the quantitative analysis of the quasidiffractive processes $\gamma\gamma \rightarrow \eta_c\eta_c$ and $\gamma\gamma \rightarrow \eta_cX(q\bar{q})$ within the three gluon exchange mechanism. The main merit of those processes is that the corresponding cross-sections can be, in principle calculated within perturbative QCD. We have estimated the corresponding cross-sections for the processes $e^+e^- \rightarrow e^+e^-\eta_c\eta_c$ and $e^+e^- \rightarrow e^+e^-\eta_cX$ with untagged e^+e^- which were found to be within the range

1—20 fb depending upon incident C.M. energy $\sqrt{s}$ and on the magnitude of odderon intercept.

Acknowledgments

L.M. is grateful to J. Czyżewski for inspiring discussions and to DESY Theory Division for hospitality. This research has been supported in part by the Polish Committee for Scientific Research grants N0 2 P03B 89 13 and 2 P30B 044 14.

References

- [1] L.N. Lipatov, in “Perturbative QCD”, ed. A.H. Mueller, World Scientific, Singapore 1989, p. 441.
- [2] L.N. Lipatov, Proceedings of the 2nd Cracow Epiphany Conference on Proton Structure, ed. M. Jeżabek and J. Kwieciński, Acta. Phys. Polon. **26** (1996) 1245.
- [3] L.N. Lipatov, preprint DESY 96 - 132, hep-ph/9610276.
- [4] E.A. Kuraev, L.N. Lipatov, V.F. Fadin, Zh. Eksp. Teor. Fiz. **72** (1977) 373; Ya.Ya. Balitzkij, L.N. Lipatov, Yad. Fiz. **28** (1978) 822; J.B. Bronzan, R.L. Sugar, Phys. Rev. **D17** (1978) 585; T. Jaroszewicz, Acta. Phys. Polon. **B11** (1980) 965.
- [5] L.V. Gribov, E.M. Levin, M.G. Ryskin, Phys. Rep. **100** (1983) 1.
- [6] J. Bartels, Nucl. Phys. **B175** (1980) 365.
- [7] J. Kwieciński, M. Praszalowicz, Phys. Lett. **B94** (1980) 413.
- [8] A. Donnachie, P.V. Landshoff, Phys. Lett. **B296** (1992) 227.
- [9] R. Janik, J. Wosiek, Phys.Rev.Lett.**79** (1997) 2935.
- [10] P. Gauron, L.N. Lipatov, B. Nicolescu, Phys. Lett. **B304** (1993) 334; Z. Phys. **C63** (1994) 253.
- [11] M.A. Braun, hep-ph/9801352.
- [12] M.G. Ryskin, Z. Phys. **C57** (1993) 89.
- [13] S. Brodsky et al., Phys. Rev. **D50** (1994) 3134.

- [14] A. Schäfer, L. Mankiewicz, O. Nachtmann, Proceedings of the Workshop “Physics at HERA”, Hamburg, October 29-30, 1991, Edited by W. Buchmüller and G. Ingelman; W. Kilian and O. Nachtman, hep-ph/9712371.
- [15] J. Czyżewski, J. Kwieciński, L. Motyka, M. Sadzikowski, Phys. Lett. **B398** (1997) 400.
- [16] R. Engel, D.Yu. Ivanov, R. Kirschner, L. Szymanowski, hep-ph/9707362.
- [17] I.F. Ginzburg, D.Yu. Ivanov, Nucl. Phys. (Proc. Suppl.) **25B** (1991) 224; I.F. Ginzburg Yad. Fiz. **56** (1993) 45.
- [18] I.F. Ginzburg, D.Yu. Ivanov. Nucl. Phys. **B388**(1992) 376.
- [19] I.F. Ginzburg, S.L. Panfil, V.G. Serbo, Nucl. Phys. **B284** (1987) 685
- [20] Report of the Working Group on “ $\gamma\gamma$ Physics”, P. Aurenche and G.A. Schuler (conveners), Proceedings of the Workshop on “Physics at LEP2” , editors: G. Altarelli, T. Sjöstrand and P. Zwirner, preprint CERN 96-01.